type of particle instrumentally or visually separable by fluorescence, color and size, with sizes of the particles ranging from $0.01\mu m$ to $6\mu m$, each antibody of the 2 to 6 antibodies is conjugated to different particles, and the ratio between the number of particles and the number of cells ranges from 0.5:1 to 20:1 in the cell suspension.

\ 2 Claim 14. (Twice Amended) Kit to detect and phenotype target cells in cell suspensions by using particles coated with antibodies/ligands directed against antigenic determinants/receptors expressed on the target cells, wherein 2 to 6 antibodies or ligands each conjugated to a particle, wherein the particle is a fluorescent or dyed particle, are incubated under gentle rotation for 5-10 minutes to 2 hours with cell suspensions containing the target cells at 0°C to 25°C, followed by an enrichment procedure, and evaluation of the target cell rosettes microscopically and/or by suitable visualizing or imaging devices, wherein the kit comprises particles conjugated to antibodies/ligands, wherein one antibody is conjugated to one type of particle instrumentally or visually separable by fluorescence, color and size, with sizes of the particles ranging from 0.01 µm to 6µm, each antibody of the 2 to 6 antibodies is conjugated to the same or different particles, and the ratio between the number of particles and the number of cells ranges from 0, 5 : 1 to 20 : 1 in the cell suspension.